

A NOTE ON QUASI-GORENSTEIN RINGS

S. H. HASSANZADEH, N. SHIRMOHAMMADI, AND H. ZAKERI

ABSTRACT. In this paper, after giving a criterion for a Noetherian local ring to be quasi-Gorenstein, we obtain some sufficient conditions for a quasi-Gorenstein ring to be Gorenstein. In the course, we provide a slight generalization of a theorem of Evans and Griffith.

1. INTRODUCTION

Let R be a Noetherian local ring with maximal ideal $\mathfrak{m}$ and residue field k . Throughout, we use $E_R(k)$ to denote the injective hull of k . Also, for a non-negative integer i , we use $H_{\mathfrak{m}}^i(R)$ to denote the i -th local cohomology module of R with respect to $\mathfrak{m}$. Following [1], we say that R is a quasi-Gorenstein ring if $H_{\mathfrak{m}}^{\dim R}(R) \cong E_R(k)$. In this paper, we investigate quasi-Gorenstein rings in terms of linkage. By the concept of linkage, we mean the complete intersection linkage. More precisely, for ideals I and J , we say that I is linked to J (written $I \sim J$) if there is an R -sequence $x_1, \dots, x_n$ in $I \cap J$ such that $I = (x_1, \dots, x_n) : J$ and $J = (x_1, \dots, x_n) : I$.

To begin, using Cohen's structure theorem in conjunction with the linkage theory, we provide a criterion which shows that the ring R is quasi-Gorenstein if and only if $\widehat{R}$, the $\mathfrak{m}$ -adic completion of R , is a certain specialization of a Gorenstein local ring. This enables us to compare (in 2.4) the Gorenstein loci and the Cohen–Macaulay loci of a quasi-Gorenstein ring. The question asking when a quasi-Gorenstein ring is Gorenstein constitutes a lot of interests. In this direction, we use the above mentioned criterion to establish the results 2.5 and 2.7 which provide some sufficient conditions under which a quasi-Gorenstein ring is Gorenstein. Finally, using 2.7 and a theorem of Serre, we provide a slight generalization of a theorem of Evans and Griffith concerning the problem to know when the residue class ring of a height two prime ideal of a certain regular local ring is complete intersection.

Date: January 1, 2004.

1991 Mathematics Subject Classification. 13C40, 13D45, 13H10.

Key words and phrases. quasi-Gorenstein ring, Linkage, Local Cohomology.

2. THE RESULTS

It is a basic fact that quasi-Gorenstein rings satisfy Serre's condition S_2 . Although this fact has already been proved, but, for the reader's convenience, in the following lemma, we provide a different proof by using local cohomology tools. Following [8], we say that a proper ideal I in a Noetherian ring is unmixed if the heights of its prime divisors are all equal.

Lemma 2.1. *Every quasi-Gorenstein ring satisfies S_2 . In particular, its zero ideal is unmixed.*

Proof. Let $(R, \mathfrak{m}, k)$ be a quasi-Gorenstein ring of dimension n . To prove that R satisfies S_2 , without loss of generality, we can assume that R is complete (cf. [3, 2.1.16]); so that, by Cohen's structure theorem, there exists a regular local ring S of dimension n' such that $R \cong S/I$ for some ideal I of S . Now, by Local Duality Theorem (cf. [2, 11.2.6]) and [3, 1.2.4], $R \cong \text{Hom}_R(E_R(k), E_R(k)) \cong \text{Hom}_R(H_{\mathfrak{m}}^n(R), E_R(k)) \cong \text{Ext}_S^{n'-n}(R, S) \cong \text{Hom}_{S/(\mathfrak{x})}(R, S/(\mathfrak{x}))$, where $\mathfrak{x}$ is a maximal S -sequence in I . Using this in conjunction with [3, 1.4.19], one can deduce that R satisfies S_2 . Now, for a prime divisor $\mathfrak{p}$ of R , we have $0 = \text{depth } R_{\mathfrak{p}} \geq \min(2, \text{ht } \mathfrak{p})$; hence the final statement holds. $\square$

In [4, 5.2], Buchsbaum and Eisenbud established relations, under certain conditions, between almost complete intersection ideals and Gorenstein ideals. Then, Schenzel, in [10], improved the above result by establishing a duality between almost complete intersection ideals and quasi-Gorenstein ideals. The next proposition, which is related to the above mentioned result, shows that one can characterize quasi-Gorenstein rings by using the concept of linkage.

In the proof of the next proposition, for a local ring $(R, \mathfrak{m})$, we use $\widehat{R}$ to denote the $\mathfrak{m}$ -adic completion of R .

Proposition 2.2. *Let S be a Gorenstein local ring and I be an ideal of S of height zero. Then S/I is quasi-Gorenstein if and only if $I \sim Sx$ for some $x \in S$.*

Proof. ($\Leftarrow$) Let $R = S/I$ and assume that $\mathfrak{m}$ is the maximal ideal of R and that $n = \dim R$. Then $R = S/0 :_S x \cong Sx = 0 :_S I \cong \text{Hom}_S(R, S)$; so that, by Local Duality Theorem, $E_R(R/\mathfrak{m}) = \text{Hom}_R(R, E_R(R/\mathfrak{m})) \cong \text{Hom}_R(\text{Hom}_S(R, S), E_R(R/\mathfrak{m})) \cong H_{\mathfrak{m}}^n(R)$. Therefore R is quasi-Gorenstein.

($\Rightarrow$) Since, by 2.1, I is unmixed, it follows from [11, 2.2] that $I \sim (0 :_S I)$. Therefore we have only to prove that $(0 :_S I)$ is a principal ideal. Using the hypothesis, we easily see that $\widehat{S}/I\widehat{S}$ is also a quasi-Gorenstein ring. Thus, applying $-\otimes_S \widehat{S}$ to the relation $I \sim (0 :_S I)$, one obtains the relation $I\widehat{S} \sim (0 :_{\widehat{S}} I\widehat{S})$. Therefore, by the

same arguments as in the proof of 2.1, we have $\widehat{S}/I\widehat{S} \cong \text{Hom}_{\widehat{S}}(\widehat{S}/I\widehat{S}, \widehat{S}) \cong 0 :_{\widehat{S}} I\widehat{S} = (0 :_S I)\widehat{S}$. Hence, by [8, Exercise 8.3], $(0 :_S I)$ is a principal ideal. $\square$

Remark 2.3. It is straightforward to see that a Noetherian local ring $(R, \mathfrak{m})$ is quasi-Gorenstein if and only if $\widehat{R}$ is so. On the other hand, according to Cohen's structure theorem, $\widehat{R}$ can be described as a residue class ring of a Gorenstein local ring with the same dimension. Therefore, 2.2 provides a criterion to justify whether a given Noetherian local ring is quasi-Gorenstein.

The next corollary shows that, for a quasi-Gorenstein ring R , the Gorenstein loci ($\text{Gor}(R)$) and the Cohen–Macaulay loci ($\text{CM}(R)$) coincide.

In the course of this paper, for a finitely generated R -module M , $r(M)$ and $\mu(M)$ will denote the type of M and the minimal number of generators of M , respectively.

Corollary 2.4. *$\text{Gor}(R) = \text{CM}(R)$ whenever R is a quasi-Gorenstein ring.*

Proof. For a prime ideal $\mathfrak{p}$ of R , one can use [8, Theorem 23.2(i)] to choose a minimal prime $\mathfrak{q}$ of the ideal $\mathfrak{p}\widehat{R}$ such that $\mathfrak{q} \cap R = \mathfrak{p}$. Then we notice that the ring $\widehat{R}_{\mathfrak{q}}/\mathfrak{p}\widehat{R}_{\mathfrak{q}}$ is of dimension 0; so that it is Cohen–Macaulay. This observation in conjunction with [3, 2.1.7 and 3.3.15] enables us to assume that R is complete. Hence, there are a Gorenstein local ring S and a zero height ideal I of S such that $R \cong S/I$. Therefore, $I \sim Sx$ for some $x \in S$ by 2.2. Now, in order to prove the assertion, it is enough to show that $\text{CM}(R) \subseteq \text{Gor}(R)$. To this end, let $\mathfrak{p} \in \text{CM}(R)$ and suppose $\mathfrak{p}' \in \text{Spec}(S)$ is such that $\mathfrak{p}' \supseteq I$ and that $\mathfrak{p}$ is the image of $\mathfrak{p}'$ under the natural homomorphism $S \longrightarrow R$. Then, $R_{\mathfrak{p}} \cong S_{\mathfrak{p}'}/IS_{\mathfrak{p}'}$ possesses the canonical module $\text{Hom}_{S_{\mathfrak{p}'}}(R_{\mathfrak{p}}, S_{\mathfrak{p}'})$ which is isomorphic to $0 :_{S_{\mathfrak{p}'}} IS_{\mathfrak{p}'} = (Sx)_{\mathfrak{p}'}$. Hence, by [3, 3.3.11], $r(R_{\mathfrak{p}}) = \mu((Sx)_{\mathfrak{p}'}) = 1$. Therefore, $R_{\mathfrak{p}}$ is a Cohen–Macaulay ring of type 1; so that $R_{\mathfrak{p}}$ is Gorenstein. $\square$

M. Hermann and N. V. Trung, in [6], present a Buchsbaum quasi-Gorenstein ring $(R, \mathfrak{m}, k)$ with $\dim R = 3$ and $H_{\mathfrak{m}}^2(R) \cong k \oplus k$ which is not Gorenstein. This example motives us to the following theorem.

Theorem 2.5. *Let $(R, \mathfrak{m})$ be a quasi-Gorenstein ring of dimension n and suppose that $H_{\mathfrak{m}}^i(R) = 0$ for all integer i with $n/2 < i < n$. Then R is Gorenstein.*

Proof. If $n \leq 2$, then, by 2.1, R is Cohen–Macaulay; hence it is Gorenstein in view of 2.4. Thus, we may suppose that $n \geq 3$. Since $\widehat{R}$ is faithfully flat over R , we have $H_{\mathfrak{m}\widehat{R}}^i(\widehat{R}) = 0$ if and only if $H_{\mathfrak{m}}^i(R) = 0$. Therefore, in view of 2.3, we may further assume that there are a Gorenstein local ring $(S, \mathfrak{n})$ and a zero height ideal I of S such that $R \cong S/I$. By 2.2, I is linked to an ideal Sx for some $x \in S$. Therefore, the assumption $H_{\mathfrak{n}}^i(S/I) = 0$ for all $n - (n - \lfloor \frac{n}{2} \rfloor) < i < n$

in conjunction with [11, 4.1] implies that S/Sx satisfies Serre's condition $S_{n-\lfloor \frac{n}{2} \rfloor}$. Now, the application of local cohomology with respect to $\mathfrak{n}$ to the exact sequence $0 \rightarrow S/I \xrightarrow{x} S \rightarrow S/Sx \rightarrow 0$ shows that $H_{\mathfrak{n}}^i(S/I) = 0$ for all $i < n - \lfloor \frac{n}{2} \rfloor + 1$. Therefore, $H_{\mathfrak{n}}^i(S/I) = 0$ for all $i < n$. Hence R is Cohen–Macaulay; so that, in view of 2.4, it is Gorenstein. $\square$

Remark 2.6. The conclusion of the above theorem fails if one of the vanishing conditions $H_{\mathfrak{m}}^i(R) = 0$ ($n/2 < i < n$) is dropped. For example, by [6, 2.2], for given integers $n \geq 5$ and $\lfloor \frac{n}{2} \rfloor < j < n$, there exists a non-Gorenstein quasi-Gorenstein ring R of dimension n such that $H_{\mathfrak{m}}^i(R) = 0$ for all integers i with $n/2 < i < n$ and $i \neq j$.

An ideal I in a local Cohen–Macaulay ring is called an almost complete intersection ideal if $\mu(I) = \text{ht } I + 1$. Evans and Griffith, in [5, 2.1], proved, over a certain regular local ring, that an unmixed almost complete intersection ideal of height two is Cohen–Macaulay. In this manner we prove the following.

Theorem 2.7. *Let S be a local ring and suppose that I is an ideal of S such that S/I is quasi-Gorenstein and that one of the following two conditions holds.*

- (i) *S is Gorenstein and I is almost complete intersection.*
- (ii) *S is regular containing a field and I is of height two.*

Then S/I is Gorenstein.

Proof. Assume (i) holds. Considering a maximal S -sequence which constitutes a part of a minimal generating set of I and passing to the residue class ring, we can assume that $I = Sy$ for some $y \in S$ and that the height of I is zero. Then, by 2.2, $I \sim Sx$ for some $x \in S$. Therefore, we have an exact sequence $0 \rightarrow S/I \xrightarrow{x} S \xrightarrow{y} S \xrightarrow{x} S \rightarrow \cdots$, which implies that S/I is an i -th module of syzygy for all $i \geq 0$. Thus S/I is Cohen–Macaulay; so that it is Gorenstein by 2.4.

Now, assume (ii) holds. One may consider a maximal S -sequence in I , pass to the residue class ring and use 2.2 to construct an ideal J of height 2 which is generated by three elements such that $I \sim J$. It now follows from [5, 2.1] that S/J is Cohen–Macaulay. Hence, by [9, 2.1], S/I is Cohen–Macaulay. Therefore, by 2.4, S/I is Gorenstein. $\square$

Remark 2.8. In [7, 1.1], Kunz, essentially, proved that an almost complete intersection ring is not quasi-Gorenstein. Then, as a corollary, he pointed out that an almost complete intersection ring is not Gorenstein. Conversely, one can employ 2.7(i), in the case where S is regular, to show that the above mentioned corollary would imply [7, 1.1].

As an application of the syzygy theorem, Evans and Griffith, in [5, 2.2], proved, over a regular local ring R which contains a field, that the residue class ring of a height two prime ideal $\mathfrak{p}$ is complete intersection whenever $\text{Ext}_R^2(R/\mathfrak{p}, R)$ is principal. (Note that this last condition is sufficient for $R/\mathfrak{p}$ to be quasi-Gorenstein.) The following corollary is a slight generalization of this result.

The result [12, Proposition 5] of Serre, which indicates that a height two Gorenstein ideal of a regular local ring is complete intersection, in conjunction with 2.7(ii) leads immediately us to the following corollary.

Corollary 2.9. *Let S be a regular local ring containing a field and suppose that I is an ideal of height two such that S/I is quasi-Gorenstein. Then I is complete intersection.*

REFERENCES

- [1] Y. Aoyama, S. Goto, *On the endomorphism ring of the canonical module*, J. Math. Kyoto Univ. **25**(1985) 21–30.
- [2] M. Brodmann, R.Y. Sharp, *Local Cohomology, An Algebraic Introduction with Geometric Application*, Cambridge University Press, Cambridge, 1998.
- [3] W. Bruns, J. Herzog, *Cohen–Macaulay Rings*, revised version, Cambridge University Press, Cambridge, 1998.
- [4] D. A. Buchsbaum, D. Eisenbud, *Algebra structures for resolutions and some structure theorems for ideals of codimension 3*, Amer. J. Math. **99**(1977) 447–485.
- [5] E. G. Evans, P. Griffith, *The syzygy problem*, Ann. of Math. **114**(1981) 323–333.
- [6] M. Hermann, N. V. Trung, *Examples of Buchsbaum quasi-Gorenstein rings*, Proc. Amer. Math. Soc. V. 117, No. 3 (1993), 619–625.
- [7] E. Kunz, *Almost complete intersections are not Gorenstein*, J. Alg. **28**(1974), 111–115.
- [8] H. Matsumura, *Commutative ring theory*, Cambridge University Press, Cambridge, 1986.
- [9] C. Peskine, L. Szpiro, *Liaison des varietes algebriques*, Invent. Math. **26**(1974), 271–302.
- [10] P. Schenzel, *A note on almost complete intersections*, Seminar Eisenbud-Singh-Vogel. Vol 2, Teubner-Texte Math., vol. 48, Teubner, Leipzig, 1982, pp. 49–54.
- [11] P. Schenzel, *Notes on Liaison and Duality*, J. Math. Kyoto Univ. **22**(1982), 485–498.
- [12] J. P. Serre, *Sur les modules projectifs*, dans Séminaire Dubreil-Pisot (Algèbre et théorie des nombres) 1960/1961, n°2.

Acknowledgment. The authors would like to thank Professor M. Chardin for helpful suggestion about Theorem 2.5. Also, the second author would like to thank Professor D. Eisenbud for kind consideration to his question about unmixed almost complete intersection ideals.

INSTITUT DE MATHÉMATIQUES, UNIVERSITÉ PARIS 6, 4, PLACE JUSSIEU, F-75252 PARIS CEDEX
05, FRANCE

E-mail address: `h.hassanzadeh@tmu.ac.ir`

FACULTY OF MATHEMATICAL SCIENCES AND COMPUTER, TARBIAT MOALLEM UNIVERSITY, 599
TALEGHANI AVE., TEHRAN 15618, IRAN

E-mail address: `ne.shirmohammadi@gmail.com`

FACULTY OF MATHEMATICAL SCIENCES AND COMPUTER, TARBIAT MOALLEM UNIVERSITY, 599
TALEGHANI AVE., TEHRAN 15618, IRAN

E-mail address: `zakeri@saba.tmu.ac.ir`